

I.F.A.S. - Univ. of Florida

U.S. DEPARTMENT OF JUSTICE  
FEDERAL BUREAU OF INVESTIGATION  
WASHINGTON, D. C. 20535

# 1976



**Agricultural Marketing Service  
U.S. DEPARTMENT OF AGRICULTURE  
Memphis, Tenn. 38122 January 14, 1977**

These reports are published bi-weekly during the harvesting season and will be summarized in a comprehensive report at the end of the crop year. A detailed description of the tests shown in this report may be found in the summary report for the previous season.<sup>1/</sup> These reports are available on request from the Standardization Section, Cotton Division, Agricultural Marketing Service, U. S. Department of Agriculture, 4841 Summer Avenue, Memphis, TN 38122.

<sup>1/</sup> Summary of Cotton Fiber and Processing Test Results, Crop of 1975,  
USDA, AMS, Cotton Division, May 1976.

## COTTON FIBER AND PROCESSING TEST RESULTS, CROP OF 1976

### Discussion of Test Results

Short staple cottons tested through January 7 from the Southwestern Area are longer, coarser and weaker than last year, according to the Cotton Division, Agricultural Marketing Service, USDA. Shirley Analyzer nonlint content is lower, while picker and card waste is higher than a year ago. Yarns spun from these samples are weaker but have fewer imperfections. Spinning potential yarn number is a little lower.

Average test results for all medium staple samples tested show slightly shorter and stronger fibers than a year ago. Shirley Analyzer nonlint content is lower, but picker and card waste is higher. Yarns spun from these samples are stronger and have fewer imperfections compared to a year ago at this time. Yarn appearance is slightly lower as is the spinning potential.

Medium staple samples tested from the Southeast show fibers to be slightly longer, more uniform, coarser and stronger than a year earlier. Shirley Analyzer nonlint content is slightly lower. Yarns spun from these samples are stronger with fewer imperfections. Average spinning potential is higher.

South Central medium staple samples tested are longer, less uniform and slightly finer than a year ago. Samples are stronger at zero gage tests. Shirley Analyzer nonlint content is lower, while picker and card waste is higher than a year ago. Yarn skein strength is slightly stronger and imperfections are fewer than a year earlier. Spinning potential is lower.

Southwestern medium staple samples tested are coarser and more uniform than a year earlier. Shirley Analyzer nonlint content is lower, while picker and card waste is higher than a year ago. Yarns have a higher appearance index and fewer imperfections.

Medium staple samples from the West have slightly shorter and weaker fibers than a year ago. Picker and card waste is higher than a year ago. Yarns spun from these samples are weaker and have lower appearance grades. Yarn imperfections are fewer. The spinning potential yarn number is lower.

Long staple samples tested through January 7 this season are longer, more uniform and coarser than a year earlier. Fiber strength is about the same as last season. Shirley Analyzer nonlint content and manufacturing waste are lower this season. Yarns spun from these samples are stronger and have fewer imperfections. The spinning potential yarn number is slightly higher than it was last season.

Long staple cottons tested from the Southeast are longer and more uniform than a year ago. Fibers are coarser and stronger. Shirley Analyzer nonlint content, picker and card waste, and comber waste are all lower than last season. Yarn quality is better than a year ago.

No additional long staple lots were received from the South Central Area during this test period.

Long staple samples from the Far West have longer, more uniform and slightly coarser fibers than in the previous season. Shirley Analyzer nonlint content and picker and card waste are higher than a year ago but comber waste is lower. Yarn strength is higher. Yarns have about the same appearance grades as last season but fewer imperfections. The spinning potential yarn number is much higher.

American Pima extra long staple samples tested through January 7 this season have about the same length, strength and mike properties as a year earlier. Picker and card waste and comber waste are both lower than last season. Yarns spun from these samples have about the same quality characteristics as a year ago.

Table 1.--Cotton: Averages of fiber and processing tests from selected gin points in the United States through January 7, 1977 <sup>1/</sup>

Staple group Area, and Crop year	Fiber test results										Processing test results												
	Lots tested	Fibrograph		Mike fine- ness	Fiber strength		S A nonlint	P & C waste	Yarn quality			Spin. Potent.											
		2.5% span	50/2.5 unif.		Pct.	Rdg.			Mpsi	G/tex	Pct.		Lbs.	Index	No.								
				Inches												Pct.	Rdg.	Mpsi	G/tex	Pct.	Lbs.	Index	No.
No.	Inches	Pct.	Rdg.	Mpsi	G/tex	Pct.	Pct.	Lbs.	Index	No.	Yarn No.												
Short Staple: Southwest 1975 1976	53	.94	45	3.6	86	21	4.0	6.7	97	107	20	42											
	54	.96	45	4.3	84	21	3.4	7.0	89	110	14	40											
Medium Staple: Southeast 1975 1976	43	1.07	44	4.2	83	22	3.4	6.3	97	97	25	53											
	48	1.08	45	4.6	85	23	3.2	6.3	106	98	20	57											
South Central 1975 1976	114	1.10	45	4.3	85	23	3.1	5.7	105	101	21	59											
	122	1.08	44	4.2	88	23	2.7	6.3	107	99	17	55											
Southwest 1975 1976	34	1.06	43	3.7	83	22	3.7	6.2	103	87	32	56											
	37	1.06	45	4.0	82	22	3.4	6.7	103	93	24	55											
West 1975 1976	68	1.12	45	4.1	92	26	2.3	5.5	124	93	23	69											
	70	1.11	45	4.2	89	25	2.2	5.8	119	89	19	66											
U.S. Average 1975 1976	259	1.09	45	4.2	86	23	3.1	5.8	108	97	23	60											
	277	1.08	45	4.2	87	24	2.8	6.2	110	95	19	58											
Significant dif- ference 2/		0.02	2	0.2	2	1	0.5	0.5	4(22s)	5	2	3											

<sup>1/</sup> Based on a limited number of samples of modal quality

<sup>2/</sup> Minimum differences considered to be significant for comparisons in this table.

Table 1.--Cotton: Averages of fiber and processing tests from selected gin points in the United States through January 7, 1977 1/ (Continued)

Staple group, Area, and Crop year	Lots	Fiber Test Results						Processing Test Results									
		Length		Mike	Strength		SA Non-lint	P&C Waste	Comber Waste	Yarn Quality					SPY		
		Span	Unif		Zero	1/8" gage				carded	combed	Appearance	Imprfctns				
														Strength		carded	combed
No.	In.	Pct.	Rdg.	Mpsi	G/tx	Pct.	Pct.	Pct.	Lbs.	Lbs.	Idx	Idx	No.	No.			
22s Carded & Combed Yarn																	
Long Staple:																	
Southeast																	
17	1.09	43	4.2	85	23	3.7	9.6	18.8	91	115	110	119	20	10			
12	1.14	45	4.4	86	25	3.1	6.7	15.8	115	136	104	116	20	8			
South Central																	
6	1.11	43	4.0	88	23	3.8	9.2	18.1	104	125	110	120	19	9			
3	1.12	42	3.7	91	26	3.4	6.8	20.3	109	137	97	103	13	10			
West																	
17	1.16	45	3.3	93	26	3.1	8.7	16.1	138	158	82	96	39	22			
5	1.19	46	3.4	87	27	4.0	9.1	13.4	144	166	84	94	35	16			
U.S. Average																	
40	1.13	44	3.8	89	24	3.5	9.2	17.6	113	135	98	109	28	15			
20	1.15	45	4.0	87	26	3.3	7.3	15.9	121	143	98	108	23	11			
50's Combed Yarn																	
Extra Long Staple:																	
West																	
9	1.47	31	3.7	104	34	2.8	7.5	18.4	67	67	111	111	2	2			
16	1.48	31	3.6	102	34	2.6	6.7	15.8	65	65	111	111	2	2			
Significant Difference 2/																	
0.02 2 0.2 2 1 0.5 0.5 0.5 4(22s) 4(22s) 5 5 2 2 3																	

1/ Based on a limited number of samples of modal quality  
2/ Minimum differences considered to be significant for comparisons in this table.

Table 2 --Cotton, American upland short staple: Quality characteristics by production areas, crop of 1976

Production Area, Classification & Sample Number				Fiber Test Results										Processing Test Results - Carded Yarns													
No	Grade	Stple	32s	Digital Fibrograph		Mike	Fiber Strength		Elon- gat'n 1/8"	S.A. Non- lint		Color Raw Stock		P & C Waste	Strength		Elongation		Appearance Index		Imprfct'ns		Spin. Poten- tial				
				2.5% span	Unif		Zero	1/8" Gage		Pct	Rdg	Mpsi	G/tex		Pct	Pct	Gra	Yel	8s or 74 tx	22sor 27 tx	8s or 74 tx	22sor 27 tx		8s or 74 tx	22sor 27 tx	No	No
SOUTHWEST AREA																											
NORTHWEST TEXAS																											
ANSON																											
2	LM LT SP	52 32	0.97	47	4.2	79	20	8.1	4.6	3	3	3	3	7.6	284	93	100	PERCENT	6.4	120	100	44	24	46			
3	LM LT SP	52 31	0.94	45	3.8	79	21	7.6	4.9	3	3	3	3	7.1	274	88	7.3	6.4	120	100	41	20	41				
BURKBURNETT																											
3	SLM LT SP	42 32	0.98	47	5.2	86	21	6.8	2.6	2	3	6.2	282	90	6.8	100	PERCENT	5.6	120	110	19	12	40				
HART																											
3	MTO SP	33 30	0.91	44	3.4	89	22	7.1	3.3	3	5	7.1	285	93	7.1	80	PERCENT	6.0	130	110	22	14	43				
LOCKNEY																											
3	LM LT SP	52 32	0.97	45	3.1	82	22	6.8	5.7	3	4	6.9	313	106	7.5	80	PERCENT	6.5	120	110	36	20	56				
LORENZO																											
2	MTO SP	33 32	1.00	44	3.1	81	22	7.7	3.0	3	5	6.1	303	100	8.1	90	PERCENT	6.9	130	110	26	11	54				
3	SLM SP	43 32	0.98	44	3.0	76	22	7.9	3.8	3	5	6.0	299	100	8.5	7.1		7.1	120	110	28	15	54				
LU88OCK																											
2	LM LT SP	52 32	1.00	44	3.1	80	22	7.4	3.3	3	5	7.3	304	99	8.3	75	PERCENT	6.9	120	90	35	19	52				
3	SLM SP	43 31	0.98	44	3.1	80	22	7.6	3.7	3	5	5.6	299	99	8.3	6.7		6.7	120	100	34	17	52				
CLNEY																											
1	MTO LT SP	32 31	0.94	46	5.2	92	20	7.3	2.8	2	3	6.1	270	87	6.6	80	PERCENT	5.6	120	110	19	9	42				
RULE																											
2	SLM LT SP	42 31	0.96	43	4.5	89	21	6.5	3.0	3	3	6.6	272	83	6.1	100	PERCENT	5.4	110	110	21	10	41				
3	SLM LT SP	42 31	0.95	44	4.4	87	21	6.3	3.4	2	3	6.8	268	87	6.4	268	87	6.4	5.5	130	100	22	10	40			
SILVERTON																											
2	LM SP	53 31	0.95	44	2.6	81	21	7.2	4.6	5	7	7.9	297	93	8.5	95	PERCENT	7.3	100	80	77	42	49				
TULIA																											
3	SLM LT SP	42 30	0.87	46	4.0	90	21	7.2	4.0	3	5	7.8	278	90	7.2	90	PERCENT	5.8	120	120	19	12	38				
VERNON																											
1	SLM LT SP	42 30	0.91	46	5.0	84	20	6.7	3.3	3	3	7.7	248	77	6.5	100	PERCENT	5.4	130	100	20	8	33				
2	SLM LT SP	42 31	0.92	45	5.1	88	20	6.4	3.0	2	3	6.1	274	88	6.3	274	88	6.3	5.6	130	110	20	10	40			
3	SLM LT SP	42 31	0.93	46	4.9	85	20	7.7	3.1	2	4	6.5	273	86	6.3	273	86	6.3	5.7	130	110	20	9	39			
OKLAHOMA																											
GRANOFIELD																											
3	SLM LT SP	42 31	0.97	46	5.3	84	22	7.0	4.5	2	3	6.8	272	91	6.8	95	PERCENT	5.9	130	110	21	11	41				
1/ Reduced from 42 because of bark																											
2/ Reduced from 33 because of bark																											
3/ Reduced from 43 because of bark																											

1/ Reduced from 42 because of bark

2/ Reduced from 33 because of bark

3/ Reduced from 43 because of bark

Table 3 --Cotton, American upland medium staple: Quality characteristics by production areas, crop of 1976

Production Area, Classification & Sample Number				Fiber Test Results										Processing Test Results - Carded Yarns									
No	Grade	Stple	32s	Digital Fibrograph		Mike	Fiber Strength		Elon-gat'n 1/8"	S.A. Non-Lint	Color Raw Stock		P & C Waste	Strength		Elongation		Appearance Index		Imprfct'ns		Spin. Potential	
				2.5% span	Unif.		Zero Gage	1/8" Gage			Gra	Yel		22s or 27 tx	50s or 12 tx	22s or 27 tx	50s or 12 tx	22s or 27 tx	50s or 12 tx	22s or 27 tx	50s or 12 tx		
SOUTH CENTRAL AREA																							
MISSISSIPPI																							
WATER VALLEY																							
4 LM	51	33	0.99	42	3.1	79	22	7.2	5.2	2	2	7.4	100	31	6.4	5.1	70	60	35	29	47		
TENNESSEE																							
BRADEN																							
4 SLM LT SP	42	33	1.03	44	4.2	80	22	7.2	3.0	2	3	6.1	101	32	6.3	4.4	100	80	17	12	50		
MCLEMORESVILLE																							
3 SLM SP	43	33	1.05	43	4.0	85	22	7.6	2.4	3	5	5.7	110	36	6.5	4.8	100	70	16	12	64		
SOUTHEAST AREA																							
ALABAMA																							
MCUNDOVILLE																							
3 LM	51	34	1.10	44	3.8	80	23	7.1	2.9	2	3	5.4	108	35	6.2	4.3	90	70	29	20	60		
PRATTVILLE																							
3 SLM	41	35	1.10	44	4.6	91	24	5.6	2.4	1	3	4.2	117	39	5.6	4.3	100	70	15	14	63		
GEORGIA																							
SHELLMAN																							
3 SLM LT SP	42	34	1.11	44	4.8	78	21	7.9	2.2	3	3	6.3	95	30	6.2	4.3	90	80	16	12	51		
SOUTHWEST AREA																							
NORTHWEST TEXAS																							
HALE CENTER																							
2 SLM SP	43	31/32	0.95	45	3.4	85	21	7.4	3.2	4	6	8.1	93	30	6.2	4.3	80	60	29	22	43		
3 LM LT SP	52	32/32	0.95	46	3.6	86	22	7.5	3.2	3	4	8.2	102	33	6.2	4.5	90	70	22	18	45		
LCCP																							
2 SLM	41	32	0.97	46	4.3	84	22	7.3	3.0	1	2	7.2	99	32	6.1	4.6	90	80	21	17	45		
LUBBOCK																							
2 SLM SP	43	31/35	1.06	39	3.0	80	23	6.6	4.2	4	6	7.5	101	34	6.4	4.6	70	60	57	52	45		
LUBBOCK																							
2 SLM SP	43	31/35	1.11	40	3.3	82	24	6.2	5.8	4	6	8.0	106	35	6.1	4.4	70	60	52	40	51		
3 LM SP	53	34/34	1.09	40	2.9	77	22	7.3	5.7	3	5	8.6	108	35	6.7	4.8	70	60	51	37	50		
*100% selected for tests, less than 100% in area																							
1/ Reduced from 33 because of bark																							
2/ Reduced from 42 because of bark																							
3/ Reduced from 43 because of bark																							

\*100% selected for tests, less than 100% in area

1/ Reduced from 33 because of bark

2/ Reduced from 42 because of bark

3/ Reduced from 43 because of bark





Table 4--Cotton, American upland long staple: Quality characteristics by production areas, crop of 1976

Production Area, Classification & Sample Number				Fiber Test Results										Processing Test Results - Carded Yarns										Spin. Potential
Sample Number		Style	32s	Digital Fibrograph		Mike	Fiber Strength		Elon- gat'n 1/8"	S.A. Non- Lint	Color Raw Stock		P & C Comber Waste	Strength		Elongation		Appearance Index		Imprfct'ns				
				2.5% span	Unif.		Zero Gage	1/8" Gage			Gra	Yel		22s or 27 tx	50s or 12 tx	22s or 27 tx	50s or 12 tx	22s or 27 tx	50s or 12 tx	22s or 27 tx	50s or 12 tx	22s or 27 tx	50s or 12 tx	
No	Grade	Name & Code		In	Pct	Rdg	Mpsi	G/tex	Pct	Pct	No	No	Pct	Lbs	Lbs	Pct	Pct	No	No	No	No			
SOUTHEAST AREA																								
ALABAMA																								
41	34			1.10	45	4.6	79	24	7.7	2.2	1	2	6.4	102	35	6.1	4.6	120	80	11	10	57		
3	SLM											*	16.6	123	43	6.9	5.3	120	110	6	5			
WEST AREA																								
NEW MEXICO																								
41	37			1.18	47	3.5	84	26	6.7	4.0	0	2	7.7	143	53	6.3	5.0	80	70	37	28	109		
2	SLM											*	12.7	165	61	6.8	5.6	90	90	20	15			
50	37			1.19	45	3.3	90	29	6.2	5.7	1	2	14.4	146	54	6.5	5.6	70	60	60	43	111		
3	LM PLUS											*	14.2	176	66	7.5	6.3	90	70	25	20			
WEST TEXAS																								
41	36			1.18	45	3.5	85	26	6.1	3.6	1	2	7.8	139	52	6.5	5.4	100	70	19	14	97		
2	SLM											*	13.7	161	59	6.9	5.4	100	90	8	8			
50	36			1.19	44	2.8	87	29	6.4	3.6	1	1	9.4	143	55	6.5	5.5	80	60	31	23	105		
3	LM PLUS											*	14.5	164	60	6.9	5.7	90	70	17	14			

\* Comber Waste and Combed Yarn Data

Table 5 --Cotton, American Pima extra long staple: Quality characteristics by production areas, crop of 1976

Production Area, Classification & Sample Number				Fiber Test Results										Processing Test Results - Combed Yarns											
Grade		Style	Array Length		Mike	Fiber Strength		Elon- gat'n 1/8"	S.A. Non- Lint	Color Raw Stock		P & C Waste	Comber Waste	Strength		Elongation		Appearance Index		Imprfct'ns					
No	Name & Code		UQL	CV		Zero Gage	1/8" Gage			Gra	Yel			50s or 12 tx	80s or 7 tx	50s or 12 tx	7 tx	50s or 12 tx	80s or 7 tx	50s or 12 tx	80s or 7 tx				
32s																									
AREA																									
WEST																									
ARIZONA																									
2	CASA GRANDE	3	46	1.58	31	3.9	105	36	7.1	2.9	3	6	6.6	100 PERCENT	15.9	65	34	5.5	4.6	110	110	1	1		
SAFFORD																									
3		4	46	1.52	31	3.9	102	36	7.9	3.4	4	5	7.0	93 PERCENT	15.5	66	33	5.5	4.5	120	110	2	1		
WENDEN																									
2		4	44	1.44	33	3.6	103	34	7.1	3.0	4	5	7.4	100 PERCENT	17.9	66	33	5.4	4.4	110	100	2	1		
NEW MEXICO																									
3	COLUMBUS	3	44	1.47	30	3.5	99	34	7.8	2.2	3	5	6.6	94 PERCENT	16.0	67	33	5.6	4.7	120	110	2	1		
WEST TEXAS																									
3	EL PASO	4	44	1.43	32	3.0	104	33	7.1	4.2	4	5	7.2	85 PERCENT	17.9	67	36	5.6	4.9	110	90	3	4		
TORNILLO																									
3		3	46	1.50	29	3.3	102	35	7.6	1.7	3	5	6.2	100 PERCENT	16.7	69	37	5.7	4.7	100	100	2	2		